



STARMEC

IVERMECTIN INJECTION FOR CATTLE

ACTIVE CONSTITUENT: 10 mg/mL IVERMECTIN

RELIABLE & ASSURED IVERMECTIN
PARASITE CONTROL



ACTIVE CONSTITUENT: 10 MG/ML IVERMECTIN

Convenient one shot Ivermectin injection for control of ivermectin-sensitive internal and external parasites in cattle. For those producers which prefer injections rather than a pour-on, Starmec Injection will treat and control internal parasites including gastrointestinal roundworms, lungworms, eyeworms, sucking lice, mites and cattle tick in cattle.

APPLICATOR

iO 20ml Injector (Drencher with conversion kit) or other variable injectors up to 20ml

WITHHOLDING PERIODS

MEAT: 28 days.
MILK: DO NOT USE in lactating cows or within 28 days of calving.
ESI: 42 days.

STORAGE

Store below 25°C (air conditioning). Store bottle in carton to protect from light.

COMPARABLE PRODUCT

Bomectin Injection.

DOSING & ADMINISTRATION

Use the contents within 24 hours of first broaching of the vial. Discard the unused portion.

The recommended dose is 1 mL STARMEC Ivermectin Injection for Cattle per 50 kg liveweight (equivalent to 200 µg ivermectin per kg liveweight) by subcutaneous injection. DO NOT administer by the intravenous or intramuscular route.

LIVEWEIGHT (KG)	DOSE VOLUME (ML)	500ML
Up to 100	2	250
101 - 150	3	166
151 - 200	4	125
201 - 250	5	100
251 - 300	6	83
301 - 350	7	71
351 - 400	8	62
401 - 450	9	55
451 - 500	10	50
501 - 550	11	45
551 - 600	12	41
601 - 650	13	38

Cattle heavier than 650 kg should be dosed at 1 mL per 50 kg. DO NOT underdose.

AVAILABLE SIZES

500mL

For more information please visit hunterriverco.com.au

Bomectin is a registered trade mark of Bayer Intellectual Property GmbH.



500ML