



RIDLEY

BARASTOC

ESSENTIAL FEEDS

**CHICK
STARTER**

COMPLETE & BALANCED
CRUMBLE



WHAT WILL IT PROVIDE MY CHICKEN?

Each pellet is a concentrate of high quality ingredients that meet the nutritional needs of developing chicks.

Contains an anticoccidial as an aid in the prevention of coccidiosis*.

Presented as a small crumble which is suitable for small beaks.


*For more information about this condition refer to the Barastoc Chook Book.

BARASTOC

ESSENTIAL FEEDS

CHICK STARTER

COMPLETE & BALANCED CRUMBLE



NUTRIENT LEVELS

TYPICAL ANALYSIS AS FED PER KG

NUTRIENT LEVELS			
Min Crude Protein	19.5%	Min Calcium	1.0%
Min Crude Fat	2.5%	Min Available Phos	0.5%
Max Crude Fibre	6.0%	Min Added Copper	8.0mg/kg
Max Added Salt	0.3%	Min Added Selenium	0.3mg/kg
		Bovatec® 20CC (Lasaloacid Sodium)	0.04%



WHAT'S IN IT?

INGREDIENTS SELECTED FROM THE FOLLOWING: Cereal Grains and By-products, Legumes and their By-products, Vegetable Protein Meals, Fats and Oils, Calcium Carbonate, Di-Calcium Phosphate, Ridley Chick Starter Vitamin and Mineral Premix, Essential Amino Acids & Enzymes, Anticoccidial - Bovatec® 20CC.



PROUDLY MADE IN AUSTRALIA
USING INGREDIENTS SOURCED
FROM AUSTRALIAN FARMERS.

EXPECTED CHICK DAILY FEED INTAKE

50g per bird per day.

FEEDING GUIDE

Ensure birds have continuous access to feed – do not restrict consumption.

Self feeders designed for poultry are a good way to ensure feed is continuously available.

Ensure that chicks have access to clean, fresh water at all times.

Hatchlings can be encouraged to start feeding by scattering feed on newspaper placed around feeders.

This feed has been formulated specifically for animals as described on the package, or the label attached to the package, and must not be used for other species of animals.

CONTAINED MEDICATION: Contains 80mg/kg Lasaloacid Sodium (as Bovatec® 20CC Lasaloacid Sodium Premix) as an aid in the prevention of coccidiosis caused by *Emeria sp* in layer replacement pullets. Bovatec® 20CC Lasaloacid Sodium Premix is a registered trademark of Zoetis Inc.

WARNING: Bovatec® 20CC Lasaloacid Sodium Premix is not registered for use in laying hens (chickens or turkeys). DO NOT feed to horses or other equids. DO NOT feed to dogs.

WITHHOLDING PERIOD: Meat - Nil. Eggs - DO NOT use in laying birds within 14 days of laying where eggs or egg products are to be used for human consumption. In this instance introduce a Barastoc layer feed 2 weeks prior to the point of lay.



SCAN THE QR CODE

to visit the Barastoc Chook Book, a comprehensive guide on rearing, feeding and keeping poultry.



we put the chicken before the egg