POISON

KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS FOR ANIMAL TREATMENT ONLY

ABATECH®

ULTRA POUR-ON ROUNDWORM, LIVER FLUKE & EXTERNAL PARASITICIDE FOR CATTLE

Active Constituents: 300 mg/mL TRICLABENDAZOLE 5 mg/mL ABAMECTIN

For the treatment and control of roundworms, liver fluke (all 3 stages) and external parasites of beef cattle.

2.5 L







PEEL BACK LABEL TO REVEAL STRAP

This product is not to be given orally or injected

DOSE RATE TABLE:

LIVEWEIGHT (KG)	DOSE VOLUME (ML)	NO. OF CATTLE TREATED
51 - 75	7.5	333
76 - 100	10	250
101 - 125	12.5	200
126 - 150	15	166
151 - 175	17.5	142
176 - 200	20	125
201 - 250	25	100
251 - 300	30	83
301 - 350	35	71
351 - 400	40	62
401 - 450	45	55
451 - 500	50	50
501 - 550	55	45
551 - 600	60	41
601 - 650	65	38

For animals heavier than 650 kg, dose at a rate of 1 mL/ 10 kg body weight.

A representative sample of animals should be weighed before treatment either with scales or weighband. Dose according to the heaviest animal by liveweight in the group (cows, bulls, calves, heifers, etc.). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

ABATECH® ULTRA POUR-ON is rainfast. Rain occurring as early as 2 hours after treatment will not affect the efficacy of the product.

A white deposit may form on the back of treated animals. This is quite normal. It is not a sign of treatment failure and should not be interfered with. Do not brush off or wash off. This deposit may last for a few days or weeks depending on weather

Temporary skin reaction may occur along the application site, especially if applied during very hot weather. This should resolve without treatment. If signs persist or worsen, contact your veterinarian.

After use clean all equipment with cooking oil. Do not use water.

General Directions:

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS **AUTHORISED UNDER APPROPRIATE** LEGISLATION

MEAT: DO NOT USE less than 49 days before slaughter for human consumption.

Reef calves born to cows that were treated with ABATECH® ULTRA POUR-ON during pregnancy and calves suckling cows that have been treated during lactation must not be slaughtered for human consumption for 49 days after treatment of the dam. MILK: DO NOT USE on cows which are producing or may in the future produce milk that may be used or processed for human consumption.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 140 days before slaughter for export. Before using this product, confirm the current ESI from The Hunter River Company Pty Limited on (03) 5820 8400 or the APVMA website (www.apvma.gov.au/residues).

Beef calves born to cows that were treated with ABATECH® ULTRA POUR-ON during pregnancy and calves suckling cows that have been treated during lactation must not be slaughtered for the export market for 140 days after treatment of the

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID INSTRUCTIONS

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If skin contact occurs, remove contaminated clothing and wash skin. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

ADDITIONAL USER SAFETY

Additional information is listed in the safety data

ENVIRONMENTAL PROTECTION

Very toxic to aquatic life. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

DISPOSAL

Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTÉR collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and

tree roots in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE

Store below 30°C (room temperature).

May form explosive peroxides. May cause an allergic skin reaction. Harmful if inhaled. Suspected of causing cancer.

Use only outdoors or in a well-ventilated area. Wash all exposed external body areas thoroughly after handling. Do not eat, drink or smoke when using this product.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and

easy to do. Continue rinsing. Store locked up.

^eAbatech is a registered trademark of The Hunter River Company.

BATCH:

EXP:



member of the macrocyclic lactone family of chemicals. It is effective against sensitive strains of internal and external parasites. It also contains Triclabendazole, a member of the benzimidazoles family of chemicals, effective against liver fluke. Resistance may develop to any chemical.

ABATECH® ULTRA POUR-ON contains Abamectin, a

For the treatment and control of adult and immature stages

Gastro-intestinal roundworms

Barber's Pole Worm (Haemonchus placei) Small Brown Stomach Worm (Ostertagia ostertagi) (including inhibited stages) Small Intestinal Worm (Cooperia oncophora, Cooperia

Stomach Hair Worm (Trichostrongylus axei) Hookworm (Bunostomum phlebotomum) Large Bowel Worm (Oesophagostomum radiatum)

Whipworm (Trichuris spp.)
Thin Necked Intestinal Worm (Nematodirus spathiaer) Intestinal Threadworm (Strongyloides papillosus)

Lungworms (Dictyocaulus viviparus)

Liver fluke (Fasciola hepatica) including adult (12 week old), immature (6 week old) and early immature (4 week old)

Sucking and biting lice
Long nosed sucking lice (Linognathus vituli) Short nosed sucking lice (Haematopinus eurysternus) Biting lice (Bovicola bovis) Little sucking lice (Solenopotes capillatus)

Aids in control of

Buffalo fly (Haematobia irritans exigua)

Cattle tick (Boophilus microplus)

(Not recommended in strategic dipping programs for cattle tick control)

DIRECTIONS FOR USE

Restraints

DO NOT USE on cows which are producing or may in the future produce milk that may be used or processed for human consumption.

Contraindications:

This product is contraindicated for use in calves under 50

Dosage and Administration:

ABATECH® ULTRA POUR-ON dose rate is 1 mL/ 10 kg body

Apply along the backline of the animal, using a standard pour-on applicator, in a strip starting between the shoulder