

Description: For the control of Spinosad-susceptible lice (Bovicola ovis), including strains resistant to Synthetic Pyrethroids (SPs), in sheep up to 7 days off-shears and sheep with long wool.

Target parasites: Bovicola ovis (Biting Lice)

Dosage & Administration: Apply iO SPINOFF POUR-ON evenly along the backline from the poll to the base of the tail using a manual or powered applicator with an iO SPINOFF POUR-ON 4-hole t-bar nozzle. Halve the dose volume and apply in 2 bands, approximately 5 cm from the mid-point of the backline on each side Applying the product to the flanks will improve the spread of iO SPINOFF POUR-ON.

DOSAGE TABLE

SHEEP CATEGORY	LIVE WEIGHT (KG)	DOSE (ML)	NO. TREATED PER 5 L PACK	NO. TREATED PER 20 L PACK
Off-shears (within 7 days after shearing)	20 - 30	15	333	1333
	31 - 40	20	250	1000
	41 - 50	25	200	800
	51 - 60	30	166	666
	61 - 70	35	142	571
	Add 2.5 mL for each 5 kg above 70 kg			
Suckling lambs (unshorn)	Up to 9	5	1000	4000
	10 - 19	7.5	666	2666
	20 - 29	15	333	1333
	Add 3.5 mL for each 5 kg above 29 kg			
Long wool (6-12 months after shearing)	Adu l t (>6 months old)	50	100	400

Withholding periods

WITHHOLDING PERIODS

MEAT: Off Shears: DO NOT USE less than 14 days before slaughter for human consumption.

Suckling Lambs: DO NOT USE less than 14 days before slaughter for human consumption.

Long Wool: Zero (0) days

MILK: DO NOT USE on ewes which are producing or may in the future produce milk that may be used or processed for human consumption.

WOOL: Zero (0) days.

Use of this product may result in wool residues that may not comply with European Union environmental quality standards. Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.

EXPORT SLAUGHTER INTERVAL (ESI)

DO NOT use less than 21 days (off shears and suckling lambs), or 0 days (long wool) before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from AIRR on (03) 5821 0922 or the APVMA website (www.apvma.qov.au/residues).

Recommended applicators: 15ml iO Applicator with 4 hole T Bar (31844)

Storage instructions: Store below 30°C (room temperature) in a dry place in tightly closed original container. Do not freeze. Do not store in direct sunlight.

Safety Directions: May irritate the skin. Will damage eyes. Avoid contact with eyes and skin. If product in eyes, wash it out immediately with water. Wash hands after use. When using the product wear elbow-length PVC gloves and goggles. After each day's use wash gloves.