

blank page

**POISON**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**Apparent** 

## MCPA 750

### SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 750 g/L MCPA  
present as the DIMETHYLAMINE SALT  
SOLVENT: 280g/L DIMETHYLACETAMIDE

**GROUP 4 HERBICIDE**

For the control of broadleaf weeds in cereals, linseed, poppies, pastures, rice, sugar cane and turf as specified in the Directions for Use table

**IMPORTANT: Read the attached booklet before use.**

APVMA Approval No: 68355/123018

**AIRR APPARENT PTY LTD**  
15/16 Princes Street, Newport NSW 2106  
P 03 5820 8400 | E [enquiries@apparentag.com.au](mailto:enquiries@apparentag.com.au) | [www.apparentag.com.au](http://www.apparentag.com.au)

**DIRECTIONS FOR USE**

**Restraints:** DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.

DO NOT apply if rain is likely within 6 hours.

**1. FIELD CROPS**

SITUATION & CROP	WEEDS CONTROLLED	STATE	RATE/ha
Wheat, Barley, Oats, Triticale	Refer to Weed Table	TAS, SA only	460mL-1.45L
		QLD only	
		NSW, ACT only	460-960mL
		VIC only	
Wheat, Barley, Triticale, Cereal Rye	Capeweed, Double Gee, Erodium, Mustard, Rapistrum, Wild Turnip, Wild Radish	WA only	600mL-1.35L
		NSW, ACT, VIC only	270mL plus 350mL diuron (500g/L flowable) 330mL plus 500mL diuron (500g/L flowable)
		NSW, ACT, VIC only	330mL plus 500mL diuron (500g/L flowable)
		SA only	230mL plus 500mL diuron (500g/L flowable)
Wheat, Barley, Oats, Triticale	Soldier Thistle Skeleton Weed	SA only	460mL plus 200mL Apparent Clopyralid 300 660mL plus 500mL Apparent Clopyralid 300
Field Peas	Prickly Lettuce ( <i>Lactuca serriola</i> )	NSW, VIC, TAS, SA only	<b>Weed Stage:</b> 4 leaf, 80mL plus 125mL diflufenican (500g/L)
			<b>Weed Stage:</b> 6 leaf, 100mL plus 150mL diflufenican (500g/L)
	Charlock (Wild Mustard) ( <i>Sinapis arvensis</i> )	NSW, VIC, TAS, SA only	<b>Weed Stage:</b> 4 leaf, 80mL plus 125mL diflufenican (500g/L)
			<b>Weed Stage:</b> 6 leaf, 100mL plus 150mL diflufenican (500g/L)
Linseed	Refer to Weed Table	QLD only	460mL-1.45L
		NSW, ACT, VIC, SA, TAS only	460-730mL
Oilseed poppies ( <i>Papaver somniferum</i> )	Fat Hen ( <i>Chenopodium album</i> ), Spear Thistle ( <i>Cirsium vulgare</i> )	NSW, TAS only	1L/ha

**CRITICAL COMMENTS**

Apply through the tillering stage after the crop plants have 5 leaves and before they commence jointing.

Apply from early tillering (4 to 5 fully emerged main stem leaves plus one or more tillers) until before the boot stage (visible swelling of the head at the top of the main stem).

Apply 460mL rate after crop has reached 5 leaf stage and weeds marked # in weed table have 3-6 leaves. Higher rates after the first node can be felt at the base of a tiller and before swelling of the head can be felt in a tiller.

Apply 460mL rate after crop has reached 2-3 leaf stage and weeds marked # in weed table have 2-4 leaves. Higher rate after the crop plants have 5 leaves to fully tillered.

Apply to crop from 5 leaf to flag leaf emergence (Zadoks 15-37).

CROP: 3-4 leaf stage (Zadoks 13-14).

CROP: 4-5 leaf stage (Zadoks 14-15).

Apply by aircraft or boom with 20-100L/ha water, when most weeds have germinated and are in 2-5 leaf stage. Crop and weeds should be dry at time of application. Some temporary yellowing of crop may occur after application. Undersown sub-clovers may be slightly retarded. DO NOT apply to undersown medic.

Apply by boom spray or aircraft when the crop is at the 3-5 leaf stage (NSW, ACT and SA only) and 5 leaf stage (VIC only). Weeds must be in the 2-5 leaf stage.

DO NOT apply to undersown medic or lucerne.

DO NOT use on Kamilaroi, Kite, Olympic or Shortim Wheat.

DO NOT use on very light sandy soils.

Apply to crop at tillering stage after the crop has 5 leaves and before the crop commences jointing.

Apply to rosettes from 5-15cm diameter when the crop is between the leaf and late

tillering growth stages. It is essential that all the Skeleton weed is emerged and minimum of 5cm diameter which is usually not before August.

This rate will provide control of Skeleton weed until harvest and will substantially reduce seed numbers the following season.

**FIELD PEAS:** Apply early post-emergence after the third node stage and before the start of flowering.

**Warning:** Field peas grown on high pH soils in the presence of free lime may be less tolerant to diflufenican (500g/L).

**APPLICATION AND WEED CONTROL:** Apply when weeds are actively growing. For optimum results apply 4 to 6 weeks post-sowing. Application beyond 8 weeks post-sowing may result in reduced levels of weed control. In most situations the rate specified for each weed size will give satisfactory control. Under certain conditions such as:

- high crop and weed density,
- late season germinations,
- abnormal weed growth (including early flowering); higher rates of product (up to the maximum rate of application specified for that weed) may be required.
- Diflufenican (500g/L) will not effectively control:
  - regrowth of suppressed weeds,
  - transplanted weeds,
  - regrowth from rhizomes or roots,
  - weeds growing under stress from previous herbicide applications.

Apply when crop is 10-15cm tall with at least 170L/ha of water.

DO NOT spray after buds appear. Some wilting effect may be seen following application but crop usually recovers quickly.

DO NOT apply by aircraft.

Use only in consultation with the poppy contracting company. Apply at least one week after petal fall after poppy seed has been set. Apply by aircraft in 30-40L of water/ha. This treatment will control late germinating weeds if they are green and at the bud or early flower stage. Spraying at this time will twist weeds below the crop canopy.

DO NOT apply in horticultural areas where there is a risk of spray or vapour drift over susceptible crops.

SITUATION & CROP	WEEDS CONTROLLED	STATE	RATE/ha
Sugar Cane	Blue Top, Chinese Burr, Flannel Weed, Gambia Pea, Bell Vine, Streaked Rattle Pod, Bindweed, Pink Convolvulus, Cupids Flower, Merremia Vine, Morning Glory	QLD only	930mL
	As above plus Fat Hen, Noogoora Burr		1.45L
Rice	Dirty Dora, Docks, Starfruit, Sedges, Buddah (Butter) Pea	All States	485mL to 970mL
Flax	Refer to Weed Table	VIC, TAS only	325mL

## 2. PASTURES, SEED CROPS & TURF

SITUATION & CROP	WEEDS CONTROLLED	STATE	RATE/ha
Grass Pastures	Refer to Weed Table	ALL STATES	460mL-2.7L
	Salvation Jane	SA only	660mL plus 400mL diuron (500g/L flowable)
	Soldier Thistle		460mL-1.45L
	Seedling and established Horehound	NSW, ACT only	2.0L
Grass Seed Crops	Refer to Weed Table	QLD, VIC, TAS, SA, WA only	460mL-2.7L
Sub-Clover Seed Crops Pastures – Spray/Graze Technique	Refer to Weed Table	VIC, SA only	460-960mL
	Calltrop, Capeweed, Charlock, Mustards, Paterson's Curse, Shepherd's Purse, Saffron, Slender Spear or Variegated Thistle, Turnip Weed, Wild Radish, Wild Turnip	NSW, ACT, VIC only	230-930mL
	Refer to Weed Table	TAS only	460mL
	Refer to Weed Table	WA only	610mL-1.25L
	Capeweed, Dandelion, Seedling Dock, Three Corner Jack, Geranium, Erodium, Mustard, Annual Thistles, Turnip Weed, Wild Radish, Wild Turnip	SA only	330mL
	Soldier Thistle		395mL
Turf	Refer to Weed Table	ALL STATES	330mL or 2.8mL per 100 sq. metres
			930mL - 1.8L or 10-19mL per 100 sq. metres

## 3. SPOT SPRAYING

SITUATION & CROP	WEEDS CONTROLLED	STATE
High Volume Spraying	Refer to Weed Table	ALL STATES
Knapsack Application	Refer to Weed Table	ALL STATES

## NON-CROP AREAS, FALLOW LAND, INDUSTRIAL & COMMERCIAL AREAS, RIGHTS-OF-WAY

SITUATION & CROP	WEEDS CONTROLLED	STATE	RATE/ha
Non-Crop areas, Fallow Land, Industrial and Commercial Areas, Rights-of-way	Refer to Weed Table	ALL STATES	Refer to Weed Table

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL  
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

CRITICAL COMMENTS
<b>Post-emergent:</b> Apply as a directed inter-row spray. DO NOT exceed rates with phenoxy sensitive varieties.
Apply by aircraft at early/mid tillering stage of rice but not later than panicle initiation. Use 485mL/ha at 35 days after aerial sowing (approx. 2 tillers per rice plant). Use 970mL/ha at 45 days or more after cereal sowing (3 tillers or more per rice plant). The crop should be sprayed before flowering of the weeds. <b>Water Management:</b> Before herbicide application, lower water level in the bays to at least two thirds of the weed growth for direct contact with MCPA 750. Weeds covered by water will not be controlled. Raise water depth to normal levels 24 hours following spraying. Apply to crop when 8-20cm high.

CRITICAL COMMENTS
Apply to established grass based pastures. Damage may occur to legumes if present. See "Legume Tolerance" below. <b>Note:</b> Capeweed, Paterson's Curse and Variegated Thistle may be poisonous to stock after spraying. Use this mixture only if spraying is delayed until early winter. Rosettes should not exceed 20cm diameter. Use low rate on seedlings with no more than 2 true leaves. Use 100 litres of water for best results. If clover present do not exceed 575mL. Apply in autumn when horehound plants are at the seedling stage. If sub-clover or medic plants are present they will be damaged by this application. Apply to established grass seed crops.
DO NOT exceed rate specified. Spray in autumn and early winter. Spray actively growing 6-8 weeks old weeds. Introduce stock 7-10 days after spraying, preferably sheep as cattle are less effective. Use stocking rate at least 5 times heavier than normal until weeds have been reduced but before survival of desirable pasture species is threatened. Lucerne and medics may be damaged and should be grazed short before spraying. See "Legume Tolerance" under General Instructions. <b>Precaution:</b> An increased quantity of poisonous plants may be eaten by stock using spray/graze e.g. Calltrop, Capeweed, Paterson's Curse and Variegated Thistle and deaths could result from causes such as nitrate poisoning. With Paterson's Curse, preferably graze stock soon destined for slaughter and avoid extended periods of grazing. Avoid grazing with young or breeding stock. DO NOT graze horses or pigs on Paterson's Curse. Observe withholding period of at least 7 days after spraying.
<b>Newly laid:</b> Delay until well established or at least two months old. Apply in high volume to actively growing weeds. See "Legume Tolerance". Buffalo Grass is more easily damaged than other grasses. <b>Established:</b> Apply in high volume to actively growing weeds. DO NOT mow for 2 days before application. Some transitory damage may occur to fine turf grasses and strawberry clover. See "Legume Tolerance". Buffalo Grass is more easily damaged than other grasses.

MIXING RATES/COMMENTS
Use 130mL/100L of water for easy to kill weeds such as Wild Turnip, Mustards and Wild Radish. Use 270mL/100L of water for more difficult to control weeds such as Docks, Hoary Cress and Bindweed. Use 13mL/10L of water for easy to kill weeds such as Wild Turnip, Mustards and Wild Radish. Use 27mL/10L of water for more difficult to control weeds such as Docks, Hoary Cress and Bindweed.

CRITICAL COMMENTS
Apply at rates shown in Weed Table when weeds only are present and non-crop or pasture is involved.

**WITHHOLDING PERIOD:****Cereal grain:** DO NOT harvest, graze or cut for stock food for 7 days after application.**Pasture:** DO NOT graze or cut for stock food for 7 days after application.**Poppies:** DO NOT harvest for 5 weeks after application.**Harvest Withholding Period (other than poppies and cereal grain):** NOT required when used as directed.**WEED TABLE:**

Where weeds are to be sprayed in a crop or pasture situation, use only the rate given for the particular crop or situation indicated under the Directions for Use table. In most cases this will give control, however some hard to kill weeds or those in advanced stages of growth may only be suppressed. The rates listed in the Weed Table below are for use where weeds only are present and no crop or pasture is involved. Use at these rates in a crop or pasture other than for spot spraying may cause damage. This product should only be used in those States where a rate or range of rates is indicated for the particular weeds listed.

APPLICATION RATE PER HECTARE							
WEEDS CONTROLLED	VIC	NSW, ACT	SA	TAS	QLD	WA	CRITICAL COMMENTS
Ball Mustard	-	-	1.15L	-	-	-	Apply at rosette stage.
Bathurst Burr	1.35L	660mL-1.35L	1.35L	1.35L	1.35L	1.35L	Spray young seedlings only.
Black Bindweed	-	970mL-1.35L	-	-	930mL-1.35L	-	Apply at young stage prior to flowering.
California Burr	-	660mL-1.35L	-	1.45L	-	-	Spray young seedlings only.
Cape Tulip	-	-	-	-	-	1.35L	Apply just prior to flowering. Gives mild suppression only.
Capeweed	-	1.45L	1.45L	1.45L	1.45L	-	Apply at seedling stage.
Carrot Weed	1.35L	-	-	-	1.35L	-	Apply at seedling stage.
Charlock	460mL	660mL	460mL	660mL-1.45L	460mL	-	Spray up to rosette stage.
Cutleaf Mignonette	-	-	1.45L	-	-	-	Apply at young rosette stage.
Dandelion	1.35L	1.35L	1.35L	1.35L	1.35L	1.35L	Apply at young rosette stage.
Deadnettle	-	1.45L	-	-	1.45L	-	Moderately susceptible. Apply at seedling stage.
Docks	960mL	-	1.45L	1.45L	-	-	Apply at rosette stage and before flower heads emerge. Only the top growth of old established plants will be affected.
Fat Hen	1.35L	660mL-1.35L	-	1.35L	1.35L	-	Spray up to rosette stage.
Fennel	1.80L	1.80L	1.80L	1.80L	1.80L	-	Apply at young stage.
Field Bindweed	1.35L	-	1.35L	1.35L	1.35L	1.35L	Apply at bud stage, repeating over several seasons.
Field Cress	-	-	-	1.45L	-	-	Apply at young stage.
Flatweed or Catsear	-	-	-	1.45L	-	930mL	Apply when weeds are 10-15cm high.
Fumitory (red)	930mL	930mL	930mL	-	930mL	-	Apply at young stage prior to flowering.
Hoary Cress	3.0L	-	1.45L	-	3.0L	-	Apply before flowering.
Horehound	-	2.0L	1.45L	-	-	-	Apply at rosette stage.
Hedge Mustard	730mL#	-	460mL	460mL	460mL	825mL	Apply at rosette stage.
Hexham Scent or Melilotus	960mL	800mL	930mL	-	930mL	-	Apply at young seedling stage.
Lesser Swinecress	-	-	-	1.45L	-	-	Apply at young stage.
Lincoln Weed	-	-	460mL	-	-	-	Apply at rosette stage.
London Rocket	-	-	-	-	-	825mL	-
Lupins	-	460mL - 960mL	-	-	-	825mL	Spray up to 10cm high.
Mintweed	1.35L	1.35L	-	-	1.35L	-	Spray young seedlings only.
Mustards	730mL#	660mL#	730mL	730mL	-	825mL	Spray up to rosette stage.
Noogoora Burr	1.35mL	660mL-1.35L	1.35L	-	1.35L	-	Spray young seedlings only.

APPLICATION RATE PER HECTARE							
WEEDS CONTROLLED	VIC	NSW, ACT	SA	TAS	QLD	WA	CRITICAL COMMENTS
Opium Poppy	-	-	-	1.45L	-	-	Apply at young stage.
Paterson's Curse	-	660-960mL	-	-	2.70L	-	Apply early rosette stage.
Pimpernel	-	-	460mL	-	-	-	Apply at rosette stage.
Plantains	-	1.35L	-	1.35L	1.35L	-	Apply at young stage.
Rapeseed	-	660-960mL	-	-	-	-	Spray up to rosette stage.
Rough Poppy	-	460-960mL	-	-	-	-	Spray up to rosette stage.
Safflower	-	460-660mL	-	-	-	-	Spray up to rosette stage.
Shepherd's Purse	-	-	-	1.45L	-	-	Apply at rosette stage.
Skeleton Weed	-	960mL-1.35L	-	-	-	-	Spray rosettes before aerial growth commences.
Sorrel	-	-	1.45L	-	-	-	Apply to rosette stage.
Stinkwort	1.35L	660-960mL	1.35L	-	-	1.35L	Apply when weeds are 10-15cm high.
Sunflower (seedlings)	-	690mL-1L	-	-	-	-	Spray up to rosette stage.
Thistles: - Artichoke	1.05L	-	1.45L	-	1.35L	460mL	Apply at rosette stage.
- Nodding	-	1.65L	-	-	-	-	Spray early rosettes. Repeat spray required in following spring or autumn.
- Saffron.	960mL-1.65L	660mL-1.35L	930mL-1.65L	930mL-1.65L	930mL-1.65L	960mL-1.65L	Spray up to rosette stage (September - WA)
- Scotch or Cotton	1.45L	-	460-930mL	1.45L	-	-	Apply at seedling stage.
- Slender	2.0L	660mL-1.35L	660mL	2.0L	-	2.0L	Spray young seedlings only.
- Soldier	730mL	-	730mL	-	-	-	Spray young seedlings. Use higher rate for larger weeds.
- Spear	-	960mL-1.35L	-	1.45L	-	-	Apply at rosette stage.
- Star	-	-	730-930mL	-	-	-	Apply at seedling stage. Use higher rate for larger weeds.
- Variegated	960mL	660mL-1.35L	460-930mL	960mL	960mL	960mL	Spray young seedlings only. Use 1.35L/ha (SA) for well-developed weeds.
Turnip Weed	730mL#	660mL#	660mL	-	-	-	Spray up to rosette stage.
Wild Artichoke	-	-	1.45L	-	-	-	Apply at rosette stage.
Wild Radish	930mL	660mL#	930mL	1.35L	460-730mL	800mL	Spray up to rosette stage.
Wild Turnip	730mL	660mL#	460mL	1.35L	460-730mL	600mL	Spray up to rosette stage.

## GENERAL INSTRUCTIONS

Weeds should be sprayed while actively growing and at their most susceptible stage. As a general guide, annuals should be sprayed when young and perennials just prior to appearance of bud stage. Extremes of cold or drought are unfavourable. The best conditions are when the soil is moist, weather fine and rain unlikely within 6 hours following application.

DO NOT spray crops or weeds outside the stages indicated in 'Critical Comments' as damage, loss of yield or inadequate weed control may result. DO NOT spray in high winds.

## APPLICATION/MIXING INFORMATION

This product mixes readily with water. Half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly before adding the remaining water.

### BOOM SPRAYING

- Use 30-120 litres of water per hectare.
- Use minimum of 170 litres water per hectare for LINSEED.

### AERIAL SPRAYING

- Use 10 - 90 litres water per hectare.

### EQUIPMENT MAINTENANCE AND USAGE

Equipment that has been used for this chemical should not be used for the application of other materials to sensitive plants unless it has been cleaned with Spraymate Tank and Equipment Cleaner or well washed out with a 2% solution of cloudy ammonia followed by several clear water rinses.

## RESISTANT WEEDS WARNING

### GROUP 4 HERBICIDE

Apparent MCPA 750 Selective Herbicide is a member of the Phenoxy group of herbicides. This product has the disruptors of plant cell growth mode of action. For weed resistance management this product is a Group 4 Herbicide.

Some naturally-occurring weed biotypes resistant to this product and other Group 4 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group 4 herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, AIRR APPARENT Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds.

### COMPATIBILITY/TANK MIXES

This product can be tank mixed with Metsulfuron 600, Banvel\*, Apparent Bromoxynil 200, Flowable Diuron, Apparent Clopyralid 300, Sertin\*, Apparent Paraquat 250, Flowable Terbutryn, Tordon\* and Tribunil\*. The effectiveness of Imidan\*, Le Mat\*, Metastox\* and dimethoate may be reduced by alkaline hydrolysis if tank mixed with this product. For these situations the addition of Spraybuff\* or Spraymate\* L1-700 is recommended. When tank mixing add the required amount of compatible product (and surfactant if recommended) when the spray tank is nearly full and while continuously agitating. If left overnight, agitate thoroughly before spraying.

### PROTECTION OF CROPS, NATIVE AND OTHER NON TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto healthy susceptible plants/crops, cropping lands or pastures.

**Drift Warning:** DO NOT use on or in situations where damage to susceptible crops – such as cotton, vines, tomatoes, vegetables, fruit trees, lucerne, legume crops and pastures, oilseed crops or other susceptible crops and trees (eg. Kurrajongs, Belahs, Eucalypts) – may occur from direct application, spray drift or vapours.

### LEGUME TOLERANCE

The use of MCPA is preferred to 2,4-D Amine (Amicide) or Ester (Estercide) in situations where legumes are undersown in the crop.

**Note:** The tolerance of pasture legumes to applied product can vary with species, variety, growing conditions, stage of growth and companion crop cover.

DO NOT spray cereal crops undersown with lucerne as damage may result.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate dams, rivers or streams with the product or used containers.

### STORAGE AND DISPOSAL

**Non-recyclable containers:** Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

**Envirodrum containers:** Store in the original sealed Envirodrum in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the drum with water or any water or any other foreign matter. After each use of the product, please ensure the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the drum have been used, please return the empty drum to the point of purchase. The drums remain the property of AIRR APPARENT Pty Ltd.

**Refillable containers:** Store in the closed original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs AIRR APPARENT Pty Ltd should be notified immediately. This minibulk/bulk container is reusable and remains the property of AIRR APPARENT Pty Ltd. DO NOT rinse empty container. Empty contents fully into application equipment. Close all valves and return to the point of supply for refill or storage. No other liquid, solid or pesticide product should be put into it. When empty return AIRR APPARENT Pty Ltd for cleaning, relabelling and refilling.

### SAFETY DIRECTIONS

Harmful if swallowed. Will damage eyes and will irritate the skin. Avoid contact with eyes and skin. When opening the container, preparing the spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. If product is on skin, immediately wash area with soap and water. If product or spray is in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126; New Zealand 0800 764 766.

### SAFETY DATA SHEET

Additional information is listed in the safety data sheet (SDS). A safety data sheet for APPARENT MCPA 750 Selective Herbicide is available from AIRR APPARENT Pty Ltd on request.

**CONDITIONS OF SALE:** AIRR APPARENT Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on AIRR APPARENT's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of AIRR APPARENT Pty Ltd has any authority to add to or alter these conditions.

\* Other trademarks