



HUNTER RIVER CO.

ABATECH®

ULTRA POUR-ON ROUNDWORM, LIVER FLUKE & EXTERNAL PARASITICIDE FOR CATTLE

ATTACKS ALL 3 STAGES OF LIVER FLUKE IN A POUR-ON!



For the treatment and control of roundworms, liver fluke (all 3 stages) and external parasites of beef cattle.

15 L

- 1L
- 2.5L
- 5L
- 15L

Active Constituent: 300 mg/mL TRICLABENDAZOLE, 5 mg/mL ABAMECTIN

Claims:

Abatech Ultra Pour-On contains triclabendazole one the most effective flukicides which kills all three stages of liver fluke including adult (12+ weeks old), immature (6-12 weeks old) and early immature (up to 6 weeks old). Treating with Abatech Ultra pour-on can have the benefit of treating an existing liver fluke population in the cattle herd, as well as preventing future pasture contamination from fluke eggs which ultimately prevents production loss caused by liver fluke disease (fasciolosis).

Abatech Ultra Pour-on also contains abamectin for a complete kill of other internal and external parasites such as sucking and biting lice which are susceptible to abamectin making Abatech Ultra Pour-on a broad spectrum endectocide and flukicide for cattle.

For the treatment and control of adult and immature stages of:

Gastro-intestinal roundworms

- Barber's Pole Worm (*Haemonchus placei*)
- Small Brown Stomach Worm (*Ostertagia ostertagi*) (including inhibited stages)
- Small Intestinal Worm (*Cooperia oncophora*, *Cooperia punctata*)
- Stomach Hair Worm (*Trichostrongylus axei*)
- Hookworm (*Bunostomum phlebotomum*)
- Large Bowel Worm (*Oesophagostomum radiatum*)
- Whipworm (*Trichuris spp.*)
- Thin Necked Intestinal Worm (*Nematodirus spathiger*)
- Intestinal Threadworm (*Strongyloides papillosus*)

Lungworms (*Dictyocaulus viviparus*)

Liver fluke (*Fasciola hepatica*) including adult (12 week old), immature (6 week old) and early immature (4 week old) larvae

Sucking and biting lice

- Long nosed sucking lice (*Linognathus vituli*)
- Short nosed sucking lice (*Haematopinus eurysternus*)
- Biting lice (*Bovicola bovis*)
- Little sucking lice (*Solenopotes capillatus*)

Aids in control of:

- Buffalo fly** (*Haematobia irritans exigua*)
- Cattle tick** (*Boophilus microplus*)

Dosage and Administration:

ABATECH® ULTRA POUR-ON dose rate is 1 mL/ 10 kg body weight. Apply along the backbone of the animal, using a standard pour-on applicator, in a strip starting between the shoulder blades. This product is not to be given orally or injected.

DOSE RATE:

LIVEWEIGHT (KG)	DOSE VOL (ML)	NO. OF CATTLE TREATED			
		1 L	2.5 L	5 L	15 L
51 - 75	7.5	133	333	666	2000
76 - 100	10	100	250	500	1500
101 - 125	12.5	80	200	400	1200
126 - 150	15	66	166	333	1000
151 - 175	17.5	57	142	285	857
176 - 200	20	50	125	250	750
201 - 250	25	40	100	200	600
251 - 300	30	33	83	166	500
301 - 350	35	28	71	142	428
351 - 400	40	25	62	125	375
401 - 450	42	22	55	111	333
451 - 500	50	20	50	100	300
501 - 550	55	18	45	90	272
551 - 600	60	16	41	83	250
601 - 650	65	15	38	76	230

WITHHOLDING PERIODS

MEAT ESI: 140 days
MILK: Do not use

For more information go to hunteriverco.com.au or contact us on (03) 5821 0922.

