

NEW



ABASTAR

Plus Selenium High Volume | ORAL DRENCH FOR SHEEP

HIGH POTENCY - SHORT ACTING!



- 1L
- 5L
- 20L

**ACTIVE CONSTITUENT: 0.8 G/L ABAMECTIN
0.4 G/L SELENIUM (AS SODIUM SELENATE)**

Claims: For the treatment and control of abamectin sensitive strains of internal parasites of sheep (including benzimidazole, levamisole and morantel resistant strains), nasal bot and itch mite of sheep and as an aid in the control of selenium deficiency in sheep.

iO ABASTAR PLUS SELENIUM contains abamectin, a member of the macrocyclic 'ML' family of drenches and also contains selenium which is an aid in control and treatment of selenium deficiency in sheep and lambs.

iO ABASTAR PLUS SELENIUM is effective against abamectin sensitive strains of the following adult and immature internal parasites of sheep:

- Adult and immature gastrointestinal roundworms:**
Haemonchus contortus – Barber's pole worm (including inhibited L4 stage)
Haemonchus placei – Large stomach worm
Ostertagia circumcincta – Small brown stomach worm (including inhibited L4 stage)
Trichostrongylus axei – Stomach hair worm
Trichostrongylus colubriformis, *Trichostrongylus vitrinus* – Black scour worm
Cooperia curticei, *Cooperia oncophora* – Small intestinal worm
Nematodirus spathiger, *Nematodirus filicollis* – Thin necked intestinal worm
Chabertia ovina – Large mouthed bowel worm
Oesophagostomum columbianum – Nodule worm
Oesophagostomum venulosum – Large bowel worm
Trichuris ovis – Whipworm
Strongyloides papillosus – Intestinal threadworm

Lungworm:
Dictyocaulus filaria – Large lungworm

Also treats and controls:
Oestrus ovis – Nasal bot
Psorergates ovis – Itch mite

Contraindications:

This product is contraindicated for use in lambs under six (6) weeks of age or less than 10 kg bodyweight.

This product is contraindicated for use when selenium intake from pasture is high or selenium is provided by other means (such as top dressing, vaccine, pellets or selenium drenches) if blood selenium levels at treatment are high. Users can determine selenium status by consulting their veterinarian.

Dosage and Administration:

iO ABASTAR PLUS SELENIUM dose rate is 1 mL/4 kg bodyweight.

DOSE RATE TABLE:

BODYWEIGHT (KG)	DOSE VOLUME (ML)	NO. OF SHEEP TREATED		
		1L	5L	10L
11 – 16	4.0	250	1,250	5,000
17 – 20	5.0	200	1,000	4,000
21 – 30	7.5	133	666	2,666
31 – 40	10.0	100	500	2,000
41 – 50	12.5	80	400	1,600
51 – 60	15.0	66	333	1,333
61 – 70	17.5	57	285	1,142
71 – 80	20.0	50	250	1,000

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 14 days before slaughter for human consumption.
MILK: DO NOT USE in ewes which are producing or may in the future produce milk that may be used or processed for human consumption.

TRADE ADVICE – Export Slaughter Interval (ESI):
 DO NOT USE less than 28 days before slaughter for export.

For SDS or technical information go to independentsown.com.au

Applicator:

iO 30ml Variable Applicator – Drench Nozzle or iO 20ml Metal Drencher.

APVMA Approval Number: 84096/118862

