



# FLYK<sup>®</sup> EXTRA

SPRAY-ON SHEEP BLOWFLY TREATMENT

65 g/L DICHLANIL

**EXTRA STRONG  
SEASON'S LONG**



**Description:** For the protection of sheep for up to 29 weeks against strike by dicyclanil-susceptible strains of blowfly (*Lucilia cuprina*). For the protection of mulesing and marking wounds on sheep against strike by dicyclanil-susceptible strains of blowflies during the wound healing process. iO FLYK Extra contains 65 g/L of dicyclanil offering premium season long protection against flystrike.

**Target parasites:** Blowflies (*Lucilia cuprina*)

**Dosage & Administration:** iO FlyK Extra is applied as a low-volume treatment on to areas where protection is required. When applying iO FlyK Extra, the nozzle of the applicator gun should be high enough above the fleece to achieve a 15 cm wide band at each pass for protection of body strike. For treatment of mulesing wounds, the band must be wide enough to cover the wound and 25 mm onto the surrounding wool.

### DOSAGE TABLE

FOR THE PREVENTION OF BODY STRIKE (DIAGRAM 1)			
WEIGHT (kg)	TOTAL VOLUME/ SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS
10-20	10	10	1
21-30	12	12	1
31-50	15	7.5	2
>50	18	9	2

FOR THE PREVENTION OF BREECH STRIKE (DIAGRAM 1)			
WEIGHT (kg)	TOTAL VOLUME/ SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS
10-20	10	10	1
21-30	12	12	1
31-50	15	7.5	2
>50	18	9	2

FOR THE PREVENTION OF FLY STRIKE AFTER MULESING (DIAGRAM 2)			
WEIGHT (kg)	TOTAL VOLUME/ SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS
Up to 30	8	4	2
31-50	10	5	2

FOR THE PREVENTION OF FLY STRIKE AFTER MARKING (DIAGRAM 3)			
WEIGHT (kg)	TOTAL VOLUME/ SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS
Ram lambs	8	4	2
Ewe lambs	4	4	1

Diagram 1.




Diagram 2.





Diagram 3.



### Withholding periods:

#### WITHHOLDING PERIODS

**MEAT:** DO NOT USE less than 14 days before slaughter for human consumption.

**MILK:** DO NOT USE on sheep which are producing or may in the future produce milk that may be used or processed for human consumption.

**WOOL:** DO NOT USE less than 3 months before shearing or fibre collection.

Use of this product may result in wool residues that may not comply with European Union environmental quality standards. Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.

#### EXPORT SLAUGHTER INTERVAL (ESI)

DO NOT USE less than 9 weeks before slaughter for export.

**Recommended applicators:** iO 30ml Drench/Pour-on applicator (25191) with spray nozzle.

**Storage instructions:** Store below 30°C (room temperature). Store in original container tightly closed in a dry, cool place. DO NOT store in direct sunlight. DO NOT re-use the container.

**Safety Directions:** Will irritate the eyes and skin. Avoid contact with eyes and skin. When using the product wear rubber gloves. Wash hands after use.