

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS
BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

io

independents Own

INCARCERATE



EASY-DOSE POUR-ON CATTLE LICE AND FLY TREATMENT

ACTIVE CONSTITUENT: 15 g/L DELTAMETHRIN

For control of Stable Fly (*Stomoxys calcitrans*), House Fly (*Musca domestica*) and Lice (*Bovicola bovis*, *Linognathus vituli* and *Haematopinus eurysternus*) and synthetic pyrethroid susceptible Buffalo Fly (*Haematobia irritans exigua*) on beef and dairy cattle.

For control of synthetic pyrethroid susceptible Buffalo Fly (*Haematobia irritans exigua*) for up to 3 weeks and for control of Stable Fly (*Stomoxys calcitrans*) for up to 1 week on horses.

Provides protection from the Biting Midge (*Culicoides brevitarsus*) for at least 14 days with effective control of blood feeding *C. brevitarsus* for 6 days post treatment.

CONTENTS

20

LITRES

★★★★★

io/10

GUARANTEE

PROUDLY



OWNED
& MADE



Made for AIRR

74-76 Drummond Rd,
Shepparton, VIC 3630.

P 03 5820 8400

independentsown.com.au

DIRECTIONS FOR USE:

Restraints: NOT TO BE USED on horses that may be slaughtered for human consumption.

Contraindications: This product is contraindicated for use on animals severely stressed or in ill health.

This product should not be used on horse's head forward of the ears.

This product should not be applied as a single spot.

Precautions: Facial skin contact may cause temporary tingling/burning.

Dosage and Administration:

DOSE RATE FOR CATTLE: 5mL per 100kg liveweight.

The following table may be used as a guide.

PESTS CONTROLLED	RATE	LIVE BODY WEIGHT	CRITICAL COMMENTS
Cattle Lice (<i>Bovicola bovis</i> , <i>Lignognathus vituli</i> , <i>Haematopinus eurysternus</i>)	5mL	Up to 100kg	Apply to animals in a race or drainage pen as a single stripe along the backline. Retreat later in the season if reinfestation occurs.
	10mL	101-200kg	
	15mL	201-300kg	
	20mL	301-400kg	
	25mL	401-500kg	
Buffalo Fly (<i>Haematobia irritans exigua</i>)	30mL	501-600kg	Apply to the whole herd – retreatment may be necessary.
	For cattle over 600kg apply 5mL per 100kg liveweight.		
Stable Fly (<i>Stomoxys calcitrans</i>)			
House Fly (<i>Musca domestica</i>)			
Biting Midge (<i>Culicoides sp.</i>)			

DOSE RATE FOR HORSES: 5mL per 100kg liveweight.

The following table may be used as a guide.

PESTS CONTROLLED	RATE	LIVE BODY WEIGHT	CRITICAL COMMENTS
Buffalo Fly (<i>Haematobia irritans exigua</i>)	5mL	Up to 100kg	Using the correct measured dose, apply to horses as two single stripes. Apply approximately 2/3 of measured dose to the midline mane area (from poll to wither) and remaining 1/3 to midline over the rump.
	10mL	101-200kg	
	15mL	201-300kg	
	20mL	301-400kg	
	25mL	401-500kg	
Stable Fly (<i>Stomoxys calcitrans</i>)	30mL	501-600kg	When applying product keep away from head forward of the ears. If product is accidentally splashed into horse's eyes, the eyes should be washed thoroughly as soon as possible, using plenty of water. Contact a veterinary surgeon for advice.
	For horses over 600kg apply 5mL per 100kg liveweight.		
			Apply to the whole herd – retreatment may be necessary.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

Horses: Treatment with iO INCARCERATE EASY-DOSE will alleviate fly annoyance related behaviour including skin flicking, head and tail twitching and feet stamping for up to 7 days. No adverse reactions were reported in horses treated with deltamethrin at 5 times the recommended dose rate.

It is recommended that the product be allowed to dry thoroughly after application before handling the animal. However, if handling or grooming the animal immediately after application is necessary wear personal protective equipment as indicated in the safety directions on this label.

General Directions:

Cattle: To avoid operator insecticide contact when concurrent treatments are carried out, iO INCARCERATE EASY-DOSE should be applied last. Operators must wear gloves.

In a proportion of treated animals, some scurfiness may develop at the site of application. In most cases this scurfiness will disappear within three weeks of application of this product.

Apply with recommended applicator.

The efficacy of this product was not affected by moderate rainfall when tested against cattle lice.

WITHHOLDING PERIODS:

CATTLE:

MEAT: Zero (0) days.

MILK: Zero (0) days.

HORSES: NOT TO BE USED on horses that may be slaughtered for human consumption.

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 21 days before slaughter for export. Before using this product, confirm the current ESI from the manufacturer on (03) 5820 8400 or the APVMA website (www.apvma.gov.au/residues).

SAFETY DIRECTIONS: Product is poisonous if swallowed. Will irritate the eyes and skin. Facial skin contact may cause temporary facial numbness. Avoid contact with eyes and skin. When using wear elbow length PVC gloves. If product on skin, immediately wash area with soap and water. After use and before eating, drinking, or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves.

FIRST AID INSTRUCTIONS: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

ADDITIONAL USER SAFETY: AVOID CONTACT WITH FOOD. Additional information is listed in the safety data sheet.

ENVIRONMENTAL PROTECTION: DANGEROUS TO FISH. DO NOT contaminate dams, rivers, streams or other waterways with the chemical or used container.

DISPOSAL: Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals onsite. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE: Store below 30°C (room temperature) in original container, tightly closed in a safe place.

APVMA APPROVAL NO.: 82675/125373

Toxic to aquatic life with long lasting effects.

Wear protective clothing and eye or face protection.

IF SWALLOWED: Rinse mouth.

Do NOT induce vomiting.

IF IN EYES: Rinse cautiously with water for several minutes.

Remove contact lenses, if present and easy to do. Continue rinsing.

Protect from sunlight.

Store in a dry and well-ventilated place. Keep cool.



B: **EXP:**